

Vaginal Delivery Care, Episiotomy Performance and Examination of Severe Perineal Tears: Cross-Sectional Study in 43 Public Hospitals

Abstract

Introduction: In recent years work has been done in the public health system in Catalonia in order to achieve consensus in promoting a model of normal childbirth care. This study aims to ascertain the proportion of singleton, full-term vaginal non-instrumental births attended in public hospitals for which an episiotomy was performed and the incidence of severe perineal tears.

Methodology: Descriptive cross-sectional study of all singleton births between 37 and 41 completed weeks of gestation attended in public hospitals (2013-2014). Births were grouped into two groups "deliveries with episiotomy" and "deliveries without episiotomy". Hospitals were grouped by level of complexity (I to III - according to the Catalonia health department classification). The relationship between qualitative variables was analysed using a chi-square test and the T-student test was used for quantitative variables; a p value equal to or lower than 0.005 was considered significant. Results: Data were collected from 53770 births. The total percentage of perineal tears was 35.7% and the total of severe tears was 0.8%. No significant differences in the mean proportion of severe perineal tears among different types of hospitals or between the two groups. The regression analysis shows that the birth weight of the newborn is the only variable that increases the risk of experiencing a severe perineal tear.

Conclusion: No significant difference was demonstrated in Catalonia in the incidence of severe perineal tears in singleton full-term vaginal non-instrumental births with and without episiotomies.

Introduction

In recent years, work has been carried out in the public health system in Catalonia (autonomous community of north east of Spain) to improve the quality of care in normal birth. The shift in the assistance model towards a less interventionist approach is based on a refocusing of maternity services that was initiated in 2007 [1] and which, in line with the Strategy for Assistance at Normal Childbirth (SANC), facilitated the progressive implementation of a more women-focused assistance model predicated upon the minimal necessary intervention.

This approach became the project for normal birth assistance, gradually introduced into healthcare practice across the region of Catalonia. This work developed via three specific courses of action aimed at improving the infrastructure in maternity wards in hospitals, raising professionals' awareness and promoting women's participation in maternity care. All of these actions are coordinated by the Catalan Ministry of Health in order to involve all stakeholders and achieve greater consensus in promoting a model of normal childbirth care that is based on known recorded services the profile of the professional best practice.

In 2013, a maternity care assessment was performed to explore obstetric interventions conducted on singleton, full-term deliveries

in all hospitals in 2014 in Catalonia [2]. In our environment care provided at birth in women with normal pregnancy is assumed mostly by midwives, but the exact proportion of births attended by midwives is not known. The assessment revealed, among other things, a downward trend in the performance of episiotomies in public hospitals and, further, no significant increase in the incidence of severe perineal tears [3]. Based on this evaluation, new lines of research were undertaken to respond to aspects of maternal and child health requiring improvement but for which the information needed for planning effective services currently does not exist. It has been shown that midwife-led care during low/medium risk pregnancy, labour, delivery and postpartum is associated with benefits for mothers and newborns, with no

Research Article

Volume 7 Issue 6 - 2017

Ramón Escuriet^{1,2*}, María J Pueyo^{1,3},
Laura García-Lausín^{3,4}, Noemí Obregón⁵,
Mercedes Pérez-Botella⁶, José M López⁷,
Isabel Salgado-Poveda⁸, María Martínez-
Pascual⁹, Pilar Asin-Chinchilla¹⁰, Mercedes
Santos-Vivas¹¹, Fátima León-Larios¹²,
Joanna White¹³ and Participants in the
MIDCONBIRTH Study

¹Directorate-General for Health Planning, Ministry of Health of the Government of Catalonia, Spain

²Mar University School of Nursing, Pompeu Fabra University, Spain

³Department of Experimental and Health Sciences, Pompeu Fabra University, Spain

⁴Mar Hospital, Mar Health Park, Spain

⁵ParcTaulí Hospital, Spain

⁶School of Community Health and Midwifery, University of Central Lancashire, UK

⁷Pallars' Regional Hospital, Spain

⁸Vall d'Hebron University Hospital, Spain

⁹Figueres Hospital, Spain

¹⁰Sexual and Reproductive Health Unit, Spain

¹¹Sant Joan de Reus Hospital, Spain

¹²University College of Nursing, Physical Therapy and Podiatry, University of Seville, Spain

¹³Department of Health and Social Sciences, University of the West of England (UWE), UK

*Corresponding author: Ramón Escuriet, Mar School of Nursing, University Pompeu Fabra, Doctor Aiguader, 80, 3^a Planta (Despatx 61.311) 08003 Barcelona Spain, Tel: +34-606995188; Email: rescuriet@gencat.cat

Received: January 31, 2017 | Published: September 11, 2017

adverse effects in comparison with models of care offered by obstetricians or when both midwives and doctors [4,5] share the work. In addition, a reduction in the use of epidural analgesia, fewer episiotomies and instrumental deliveries have been observed during midwife-led care in some settings [6,7].

One of the informational needs identified in the assessment performed in 2014 was the proportion and the outcomes of normal births attended to in public hospitals by midwives, and to deepen the results of the episiotomy performance. With the aim to obtain all this new information the study MidconBirth(MCB) was initiated [8].

This work explores existing data in hospital discharge records with the aim to identify the performance of episiotomy of low and medium risk births in a specific period of time. It is important to know for the understanding of this study that all hospitals do attend low and medium risk births regardless its level of complexity. Women at low or medium risk are then treated under the same guidelines in all hospitals, so results may be compared between hospitals grouped by levels of complexity.

The procedure of episiotomy– an intervention that can be performed by midwives during vaginal non-instrumental delivery – was chosen for this analysis as a sensitive indicator, together with the occurrence of severe perineal tear, which could be related to the performance or non-performance of this intervention. The findings presented show the overall rate of episiotomies and perineal tears recorded in all public hospitals in the region of Catalonia, without differentiating the type of professional attending each childbirth. This baseline information will be useful

in comparing the results obtained in the subsequent study (MCB), which will collect data on these same indicators, disaggregated in relation to births attended by midwives and/or other health professionals.

Ethical approval

This study was exempt from the ministry of health of the government of Catalonia ethics committee approval as is used publicly available anonymized data.

Methodology

The descriptive cross-sectional study of births recorded in the Minimum Basic Data Set (MBDS) in Catalonia), during 2013-2014 was used. The register is mandatory for all 43 public hospitals and provides the basis for government reimbursement through the Catalan Health Service. The International Classification of Diseases (ICD-9-CM) is used for coding in the MBDS.

Data on all singleton births between 37 and 41 completed weeks of gestation attended in public hospitals were retrieved (Figure 1). Subsequently, cases without associated codes indicating an instrumental delivery or caesarean section were selected. Once cases were compiled, they were grouped into two categories “deliveries with episiotomy” and “deliveries without episiotomy” as well as according to codes representing the type of perineal tear registered: 1st degree tear, 2nd degree tear, severe perineal tear (groups 3 and 4) and “other tears” (where tears were reported but with no specified degree). In addition, information on maternal age at delivery and newborn birth weight was collected for each case.




Figure 1: Flow-Chart.

Due to the small number of cases of severe perineal injury recorded in a single year, for the analysis, accumulated cases from 2013 and 2014 were obtained.

A descriptive analysis was prepared for each hospital group to obtain the mean and the confidence interval (95%) for each variable. Hospitals were grouped by level of complexity (I to III),

with Level I having the lowest capacity and Level III the greatest capacity and technical resources for complex care. Figure 2 (levels of complexity). The relationship between qualitative variables was analysed using a chi-square test and the T-student test was used for quantitative variables; a *p* value equal to or lower than 0.005 was considered significant. Finally, a logistic regression model was used to analyse the relationship between maternal

age, birth weight and performance of an episiotomy and severe perineal tear. The PASSW 21 statistical package was used for the analysis.

Results

Data were collected from 53770 births (26697 from 2013 and 27073 from 2014) Figure 1. Combined cases from 2013 and 2014 were used for the bivariate analysis and logistic regression cases. There were a total of 53770 SFTVNIB births, of which 77.10% had no episiotomy and 22.90% had an episiotomy. The percentage of severe perineal tears recorded in all births was 0.80% (0.7 to 0.85%). The average birth weight at birth was 3330.1 g, (SD 434.8, range 1000.0-5620.0). The average maternal age was 30.7 years (SD 5.6) (minimum maternal age at birth was 12 years – and the maximum maternal age was 51 years). The classification of hospitals is shown in Table 1; 32.9% of births were attended in level I hospitals; 31.7% in level II and 35.3% in level III hospitals.

Figure 2 shows the perineal tears recorded in all singleton full-term vaginal non-instrumental births (SFTVNIB) attended in Catalonia throughout 2013-2014. Births are classified into two groups “deliveries without episiotomy” and “deliveries with episiotomy” and in each group the mean proportion of the overall number of tears recorded in Catalonia are shown.

To explore the possible differences between types of hospital, tears registered in the two groups of births were explored in relation to the type of hospital in which they occurred and were

treated. Table 1 shows the mean ratio and confidence interval (95%) of each type of tear in each group of hospitals and for each group (births with episiotomy and births without episiotomy). Figures obtained show no significant differences in the mean proportion of severe perineal tears among different types of hospitals or between the two groups. However, a relevant difference is seen for 1st and 2nd degree tears between the two groups of births in all groups of hospitals. The highest mean proportion of 1st degree perineal tears is recorded in births without an episiotomy and attended in level I hospitals.




Figure 2: Perineal tears in Singleton Full Term Vaginal Non Instrumental Births (SFTVNIB).

Table 1: Hospitals classification.

Hospital Complexity Level	Number of Hospitals	Hospital Capacity for Complexity Care	
Level I	23	Low and medium risk term births, premature births [35-36 weeks gestation]. In case of complications stabilisation and transfer	
Level II	Level IIA	6	High risk deliveries not requiring other medical specialties, premature births (> 32 weeks gestation or >1.5 Kgrs). Neonates with non severe pathology, mechanical ventilation for short periods and post NICU care. Providing support to Level I hospitals
	Level IIB	5	Very high risk births. Intermediate obstetric risk units including shorttime mechanical ventilation. Premature births (>28 weeks gestation or 1.0 Kgrs). Congenital anomalies nonrequiring specialist. Minorsurgical problems. Providing support to Level I, IIA
Level III	Level IIIA	6	Pregnancies and deliveries requiring ongoing coordination with other medical specialties. Extreme prematures (<28 weeks of gestation or < 1.0 Kgrs), Providing support to Level I, IIA and IIB
	Level IIIB	2	Deliveries requiring the highest level of medical and surgical complexity. Permanent coordination with sub-specialties (fetaal heart surery, transplantation, etc.). Providinig support to Level I, IIA, IIB and IIIA
Level of Neonatal Care I: Health Newborn>35-36 weeks of Pregnancy; Level of Neonatal Care IIA: newborn>32 weeks of Pregnancy and/or >1000g; Level of Neonatal Care IIB: newborn>28 weeks of Pregnancy and/or >1000g; Level of Neonatal Care IIIA: newborn< 28 weeks of Pregnancy and/or 1000g; Level of Neonatal Care IIIB: Cardiac Surgery Neonates, Neurosurgery			

The bivariate analysis shows that episiotomies were performed more frequently on younger women (mean age 29.99 years, SD 5.76); this difference was significant ($p = 0.000$) compared to the mean age (30.99 years SD 5.52) of women who did not have an episiotomy. No significant difference ($p = 0.977$) was found in the age of women with severe perineal tears.

With regard to infant weight at birth, the bivariate analysis revealed a significant difference in the weight of newborns between births in which an episiotomy was performed (greater mean weight), and those in which an episiotomy was not performed ($p = 0.009$), although it was deemed that the weight difference identified (11 g) had no clinical relevance. However, a significant

clinically relevant difference was found in the weight of newborns in which a severe tear occurred (average weight 3489.29 g SD 441.72) and birth weight of newborns whose mothers did not have a severe perineal tear (mean weight 3330.24 g, SD 434.53 g). Finally, the difference between severe perineal tears among births with or without an episiotomy was not found to be significant ($p = 0.459$)

For the regression analysis, a severe perineal tear was considered a variable of the result and maternal age, birth weight and episiotomy as explanatory variables. The analysis shows that the birth weight of the newborn is the only variable that increases the risk of experiencing a severe perineal tear (Table 2 & 3).

Table 2: Perineal tears (SFTVNIB) according to type of hospital.

			%	CI
No Episiotomy	H. Level I	Perineal Tear 1º	33.07	32.3-33.9
		Perineal Tear 2º	12.83	12.30-13.40
		Severe Perineal Tear	0.65	0.50-0.80
		Other Perineal Tear	1.18	1.00-1.40
	H. Level II	Perineal Tear 1º	27.76	27.00-28.60
		Perineal Tear 2º	13.46	12.90-14.10
		Severe perineal Tear	0.70	0.60-0.80
		Other perineal Tear	1.13	0.90-1.30
	H. Level III	Perineal Tear 1º	25.23	24.50-25.90
		Perineal Tear 2º	12.10	11.60-12.30
		Severe Perineal Tear	0.98	0.80-1.10
		Other Perineal Tear	1.72	1.50-1.90
Episiotomy	H. Level I	Perineal Tear 1º	6.64	5.90-7.40
		Perineal Tear 2º	2.80	2.30-3.30
		Severe Perineal Tear	0.75	0.50-1.00
		Other Perineal Tear	0.36	0.20-0.50
	H. Level II	Perineal Tear 1º	2.47	2.00-2.90
		Perineal Tear 2º	2.1	1.70-2.50
		Severe Perineal Tear	0.61	0.40-0.80
		Other Perineal Tear	0.22	0.10-0.40
	H. Level III	Perineal Tear 1º	7.48	6.60-8.40
		Perineal Tear 2º	4.30	3.60-5.00
		Severe Perineal Tear	0.8	0.50-1.10
		Other Perineal Tear	1.15	0.80-1.50

Table 3: Regression analysis. Perineal tears and risk factors.

Maternal Age			Newborn Weigh at Birth			Episiotomy		
OR	CI (95%)	p value	OR	CI (95%)	p value	OR	CI (95%)	p value
0.997	0.980-1.015	0.752	1,001	1.001-1.001	0.000	1,083	0.857-1.370	0.503

Discussion

The current study was conducted using data available in the MBDS from which a sample of perineal tears in singleton full-term vaginal non-instrumental births was selected. The analysis of 53770 births included for the years 2013 and 2014 revealed that the proportion of episiotomies performed was 22.9%, showing a reduction in relation to the analysis performed using the same methodology in 2007, which found a 29.2% ratio of episiotomies in singleton full-term vaginal non-instrumental births [2]. Data collected demonstrate that, in Catalonia, the systematic practice of performing episiotomies in this particular type of births has decreased and is now carried out more restrictively, as prescribed by practice-based evidence [9].

Completeness and accuracy of the records of episiotomies and perineal tears was verified via a comparison of existing data in hospitals and data recorded in the MBDS by means of physical visits to 30 public hospitals throughout 2012-2013 [10]. This finding, together with the standardization of the records and examination of the diagnoses and related procedures recorded in each case, represent one of the strengths of this study. By examining data that have been recorded in a standardized manner for several years as part of the same registration system, we have biases already described by other authors who attribute the increase in severe tears recorded in recent years to improved registration and standardized classification [11-15]. The use of separately coded data also helped strengthen the study's findings.

Our study presents the proportions of each type of perineal tear recorded in 2014, in both births in which an episiotomy was performed as well as in those without an episiotomy. The overall results of this study coincide with other studies [16-18] that demonstrate a higher proportion of 1st and 2nd degree tears in births without an episiotomy, while it also confirms that the proportion of severe perineal tears remain lower than 1% of births in which an episiotomy was performed as well as in those without an episiotomy. The proportion of severe perineal tears found in Catalonia is lower than that found in the majority of studies conducted [19-21] probably because, in this study, instrumental deliveries were excluded, as they usually exhibit a higher incidence of perineal tears. The presentation of data for each group of hospitals according to their level of complexity makes it possible to observe that the proportion of severe perineal tears of less than 1% remained constant for the three groups of hospitals. Level I hospitals with low and medium risk term births, midwives being in charge of the care of the woman and childbirth and in case of complication stabilisations and transfer to a higher level hospital is required. Level II, hospitals with high risk deliveries not requiring other specialities, premature births up to 28 weeks gestation, and level III hospitals, pregnancies and

deliveries requiring ongoing coordination with other medical specialities, and deliveries requiring the highest level of medical and surgical complexity and extreme prematures (less than 25 weeks gestation or less than 1kg.)

Our study was conducted in public hospital settings in a region in which a care model based on normal delivery with minimal intervention is being implemented. The non-routine practice of episiotomies, among other recommendations, has been promoted within this care framework. Our assessment confirms, as found in other studies, that 1st degree tears, considered benign lesions that do not require repair, are the most frequent occurrence in births where an episiotomy has not been performed [22-24] and that 2nd degree tears are also more frequent among births without episiotomies, but the restrictive use of an episiotomy has prevented a greater proportion of women having had a surgical incision as opposed to a minor tear [25]. Finally, the study revealed how the restrictive use of episiotomies reduces their percentage without compromising the safety of women or newborns [26,27].

Various studies have revealed several risk factors for severe perineal tears. The most frequently reported risks are parity, maternal age, birth weight of the newborn and the use of instruments, but also maternal position during labour, use of epidural analgesia, induction of labour, stimulation with oxytocin and sex of the child [18,28-30]. Other, earlier studies served to rule out risk factors related to the occurrence of perineal tear, such as hands-off, use of a bathtub or even maternal position during labour [30,31] or identified protective factors against severe lacerations such as lateral position [32].

This study was conducted as part of the assessment of the Normal Birth Care Programme by the Ministry of Health of the Government of Catalonia, which recommends not performing episiotomies as a matter of routine, as routine episiotomy was the practice before introducing the programme. After six years of implementing the Programme, the results from this study confirm that the practice of episiotomies can be reduced without causing injury [33] and that factors related to the type of delivery care that can increase the likelihood of perineal tears should continue to be explored.

Contribution of the study

The work that we present in this paper, analyses the situation with the current available information in the MBDS, prior to the start of a new study (MCB). Results coming from this work are important for further comparisons with upcoming studies if it is demonstrated, in future works that there are other different variables, that may contribute with new information to explain perineal tear rates and to permit a broader understanding of birth outcomes

Limitations

This study was not able to analyse parity because this indicator was not available in the database.

Conclusion

No significant difference was demonstrated in Catalonia in the incidence of perineal tears in singleton full-term vaginal non-instrumental births with and without episiotomies. Non-performance of an episiotomy in these births did not increase the likelihood of experiencing a serious tear.

Acknowledgement

This study is part of the MidconBirth Project, comprising members of Working Group 4 of the ISCH COST 1405. This study contributes to ISCH COST Action 1405 Building Intrapartum Research through Health – an interdisciplinary whole system approach to understanding and contextualising physiological labour and birth (BIRTH).

References

1. Ministry of Health and consumer's affaires (2007). Strategy for Assistance at Normal Childbirth in the National Health System. Madrid, Spain.
2. Escuriet R, Pueyo MJ, Biescas H, Espiga I, White J, et al. (2014) Obstetric interventions in two groups of hospitals in Catalonia: a cross-sectional study. *BMC Pregnancy Childbirth* (14): 143.
3. Escuriet R, Pueyo MJ, Perez-Botella M, Espada X, Salgado I, et al. (2015) Cross-sectional study comparing public and private hospitals in Catalonia: is the practice of routine episiotomy changing? *BMC Health Serv Res* 15: 95.
4. Sandall J, Soltani H, Gates S, Shennan A, Devane D (2013) Midwife-led continuity models versus others models of care for childbearing women. *Cochrane Database Syst Rev* 4: CD004667.
5. Kenny C, Devane D, Normand C, Clarke M, Howard A, et al. (2015) A cost-comparison of midwife-led compared with consultant-led maternity care in Ireland (the MidU study). *Midwifery* 31(11): 1032-1038.
6. Miller S, Abalos E, Chamillard M, Ciapponi A, Colaci D, et al. (2016) Beyond too little, too late and too much, too soon: a pathway towards evidence-based, respectful maternity care worldwide. *The Lancet* 388(10056): 2176-2192.
7. Escuriet R (2015) MIDCONBIRTH-Midwives' contribution to normal childbirth care: Cross-sectional study in public hospitals in Catalonia (ISRCTN14062994 DOI 10.1186/ISRCTN14062994).
8. MSSSI (Ministerio de Sanidad, Servicios Sociales e Igualdad), Guía práctica clínica sobre la atención al parto normal, Febrero 2011.
9. Escuriet R, Pueyo M, Biescas H, Colls C, Espiga I, et al. (2014) Obstetric interventions in two groups of hospitals in Catalonia: a cross-sectional study. *BMC Pregnancy Childbirth* 14: 143.
10. Escuriet R, Modelos de organización de los servicios de atención al parto: efectos sobre la provisión de servicios y los resultados, Tesis Doctoral, Noviembre 2015.
11. Gurol-Urganci I, Cromwell DA, Edozien LC, Mahmood TA, Adams EJ, et al. (2013) Third and fourth degree perineal tears among primiparous women in England between 2000 and 2012: time trends and risk factors. *BJOG* 120(12): 1516-1525.
12. Ampt AJ, Ford JB, Roberts CL, Morris JM (2013) Trends in obstetric anal sphincter injuries and associated risk factors for vaginal singleton term births in New South Wales 2001–2009. *Aust N Z J Obstet Gynaecol* 53(1): 9-16.
13. McLeod NL, Gilmour DT, Joseph KS, Farrell SA, Luther ER (2003) Trends in major risk factors for anal sphincter lacerations: a 10-year study. *J Obstet Gynaecol Can* 25(7): 586-593.
14. Burns EM, Rigby E, Mamidanna R, Bottle A, Aylin P, et al. (2012) Systematic review of discharge coding accuracy. *J Public Health* 34(1): 138-148.
15. Viswanathan M, Hartmann K, Palmieri R, Lux L, Swinson T, et al. (2005) The use of episiotomy in obstetrical care: A systematic review. Number 112. AHRQ Publications Number 05-E009-1, Agency for Health Care research and quality, Rockville, MD, USA.
16. Klein MC, Gauthier RJ, Jorgensen SH, Robbins JM, Kaczorowski J, et al. (1992) Does episiotomy prevent perineal trauma and pelvic floor relaxation? *Online J Curr Clin Trials Doc* No 10.
17. Eltorkey MM, Nuaim MA (1994) Episiotomy, elective or selective: A report of a random allocation trial. *J Obstet Gynaecol* 14(5): 317-320.
18. Penneycard JM, Cunningham Y, Love C (2012) Third and fourth degree perineal tear audit, NHS Lothian. *Archives of Disease in Childhood Fetal & Neonatal* 97(Supplement 1): A101-A102.
19. Eskandar O, Shet D (2009) Risk factors for 3rd and 4th degree perineal tear. *J Obstet Gynaecol* 29(2): 119-122.
20. Hudelist G, Gelle'n J, Singer C, Ruecklinger E, Czerwenka K, et al. (2005) Factors predicting severe perineal trauma during childbirth: role of forceps delivery routinely combined with mediolateral episiotomy. *Am J Obstet Gynecol* 192(3): 875-881.
21. Chehab M, Courjon M, Eckman-Lacroix A, Ramanah R, Maillet R, et al. (2014) [Impact of a major decrease in the use of episiotomy on perineal tears in a level III maternity ward]. *Jo J Gynecol Obstet Biol Reprod (Paris)* 43(6): 463-469.
22. da Silva FM, de Oliveira SM, Bick D, Osava RH, Tuesta EF, Riesco ML (2012) Risk factors for birth-related perineal trauma: a cross-sectional study in a birth centre. *J Clin Nurs* 21(15-16): 2209-2218.
23. Gerdin E, Sverrisdottir G, Badi A, Carlsson B, Graf W (2007) The role of maternal age and episiotomy in the risk of anal sphincter tears during childbirth. *Aust N Z J Obstet Gynaecol* 47(4): 286-290.
24. Harvey MA, Pierce M, Alter JE, Chou Q, Diamond P, et al. (2015) Obstetrical Anal Sphincter Injuries (OASIS): Prevention, Recognition, and Repair. *J Obstet Gynaecol Can* 37(12): 1131-1148.
25. Lai CY, Cheung HW, Hsi Lao TT, Lau TK, Leung TY (2009) Is the policy of restrictive episiotomy generalisable? A prospective observational study. *J Matern Fetal Neonatal Med* 22(12): 1116-1121.
26. Dannecker C, Hillemanns P, Strauss A, Hasbargen U, Hepp H, et al. (2004) Episiotomy and perineal tears presumed to be imminent: randomized controlled trial. *Acta Obstet Gynecol Scand* 83(4): 364-368.
27. Elvander C, Ahlberg M, Thies-Lagergren L, Cnattingius S, Stephansson O (2015) Birth position and obstetric anal sphincter injury: a population-based study of 113 000 spontaneous births. *BMC Pregnancy Childbirth* 15: 252.
28. Pergialiotis V, Vlachos D, Protopapas A, Pappa K, Vlachos G (2014) Risk factors for severe perineal lacerations during childbirth. *Int J Gynaecol Obstet* 125(1): 6-14.

29. Dahlen H, Priddis H, Schmied V, Sneddon A, Kettle C, et al. (2013) Trends and risk factors for severe perineal trauma during childbirth in New South Wales between 2000 and 2008: a population-based data study. *BMJ Open* 3(5).
30. Smith LA, Price N, Simonite V, Burns EE (2013) Incidence of and risk factors for perineal trauma: a prospective observational study. *BMC Pregnancy Childbirth* 13: 59.
31. De Jonge A, Van Diem MT, Scheepers PL, Buitendijk SE, Lagro-Janssen AL (2010) Risk of perineal damage is not a reason to discourage a sitting birthing position: a secondary analysis. *Int J Clin Pract* 64(5): 611-618.
32. Prinyankur R, Sujatha MS, Bivas B, Anumita C, Pijushkanti R (2015) A comparative study of perineal morbidity in vaginal delivery with and without episiotomy. *Int J Reprod Contracept Obstet Gynecol* 4(5): 1442-1445.
33. Räisänen S, Vehviläinen-Julkunen K, Heinonen S (2010) Need for and consequences of episiotomy in vaginal birth: a critical approach. *Midwifery* 26(3): 348-356.